arXiv:1404.7663v1 [math.OA] 30 Apr 2014

THE JIANG-SU ABSORPTION FOR INCLUSIONS OF UNITAL C*-ALGEBRAS

HIROYUKI OSAKA* AND TAMOTSU TERUYA

Abstract. We introduce the tracial Rokhlin property for a conditional expectation for an inclusion of unital C*-algebras $P \subset A$ with index finite, and show that an action α from a finite group G on a simple unital C*-algebra A has the tracial Rokhlin property in the sense of N. C. Phillips if and only if the canonical conditional expectation $E\colon A \to A^G$ has the tracial Rokhlin property. Let $\mathcal{C}$ be a class of infinite dimensional stably finite separable unital C*-algebras which is closed under the following conditions:

(1) $A \in \mathcal{C}$ and $B \cong A$, then $B \in \mathcal{C}$.
(2) If $A \in \mathcal{C}$ and $n \in \mathbb{N}$, then $M_n(A) \in \mathcal{C}$.
(3) If $A \in \mathcal{C}$ and $p \in A$ is a nonzero projection, then $pAp \in \mathcal{C}$.

We prove that for any simple unital C*-algebra A which is also local tracial $\mathcal{C}$-algebra in the sense of Fan and Fang, if any C*-algebra in $\mathcal{C}$ is weakly semiprojective and $E\colon A \to P$ is of finite index type with the tracial Rokhlin property, P is a unital local tracially $\mathcal{C}$-algebra. The main result is that if A is simple, separable, unital nuclear, the Jiang-Su absorbing and $E\colon A \to P$ has the trcial Rokhlin poperty, then P is the Jiang-Su absorbing. As an application, when an action α from a finite group G on a simple unital C*-algebra A has the tracial Rokhlin property, then for any subgroup H of G the fixed point algebra A^H and the crossed product algebra $A \rtimes_{\alpha_{|H}} H$ has the Jiang-Su absorbing. We also show that the strictly comparision property for a Cuntz semigroup $W(A)$ is hereditary to $W(P)$ if A is simple, separable, exact, unital, and $E\colon A \to P$ has the trcial Rokhlin property.

1. Introduction

The purpose of this paper is to introduce the tracial Rokhlin property for an inclusion of separable simple unital C*-algebras $P \subset A$ with finite index in the sense of [33], and prove theorems of the following type. Suppose that A belongs to a class of C*-algebras characterized by some structural property, such as tracial rank zero in the sense of [14]. Then P belongs to the same class. The classes we consider include:

- Simple C*-algebras with real rank zero or stable rank one.
- Simple C*-algebras with tracial rank zero or tracial rank less than or equal to one.
- Simple C*-algebras with the Jiang-Su algebra $\mathcal{Z}$ absorption.
- Simple C*-algebras for which the order on projections is determined by traces.

Date: 28 April, 2014.
2000 *Mathematics Subject Classification.* Primary 46L55; Secandary 46L35.
Key words and phrases. Jiang-Su absorption, Inclusion of C*-algebras, strictly comparison.
*Research of the first author partially supported by the JSPS grant for Scientific Research No.23540256.

- Simple C*-algebras with the strict comparison property for the Cuntz semigroup.

The 3rd condition and 5th condition are important properties related to Toms and Winter's conjecture, that is, the properties of strict comparison, finite nuclear dimension, and Z-absorption are equivalent for separable simple infinite-dimensional nuclear unital C*-algebras ([31] and [35]).

We show that an action α from a finite group G on a simple unital C*-algebra A has the tracial Rokhlin property in the sense of [24] if and only if the canonical conditional expectation $E\colon A \to A^G$ has the tracial Rokhlin property for an inclusion $A^G \subset A$. When an action α from a finite group on a (not necessarily simple) unital C*-algebra has the Rokhlin property in the sense of [9], all of the above results are proved in [22][23].

The essential observation is already done in the proof of Theorem 2.2 of [24] the crossed product $A \rtimes_\alpha G$ ($= C^*(G, A, \alpha)$ in [24]) has a local approximation property by C*-algebras stably isomorphic to homomorphic images of A. Since the Jiang-Su algebra $\mathcal{Z}$ belongs to classes of direct limits of semiprojective building blocks in [10], technical difficulties arises because we must treat arbitrary homomorphic images in the approximation property. (Homomorphic images of semiprojective C*-algebras need not be semiprojective.) In [36] they introduced the unital local tracial $\mathcal{C}$ property, which is generalized one of a local $\mathcal{C}$ property in [20], for proving that a C*-algebra with a local approximation property by homomorphic images of a suitable class $\mathcal{C}$ of semiprojective C*-algebras can be written as a direct limit of algebras in the class. When each homomorphism is injective, the unital local tracial $\mathcal{C}$ property is equivalent to the tracial approximation property in [4]. Note that when an action α from a finite group G on a simple unital C*-algebra A, the tracial approximation property for A is inherited to the crossed product algebra $A \rtimes_\alpha G$. (See [36] and Theorem 3.3.)

We know of several results like those above for tracial approximation in the literature: for stable rank one (Theorem 4.3 of [4] and [6]), for real rank zero ([6]), for the $\mathcal{Z}$-absorption (Corollary 5.7 of [8]), for the Blackadar comparson property (Theorem 4.12 in [4]).

The paper is organized as follows. In section 2 we introduce the notion of a unital local tracial $\mathcal{C}$-algebra and a tracial approximation class (TA$\mathcal{C}$ class) and we show in section 3 that when an action α from a finite group G on a simple unital C*-algebra A has the tracial Rokhlin property, then the crossed product algebra $A \rtimes_\alpha G$ belongs to the class TA$\mathcal{C}$ for A in TA$\mathcal{C}$. In section 4 we introduce the tracial Rokhlin property for an inclusion $P \subset A$ of unital C*-algebras and show that if A is a simple local tracial $\mathcal{C}$-algebra, then so is P (Theorem 4.11). In particular, if A has tracial topological rank zero (resp. less than or equal to one), so does P (Corollary 4.12). In section 5 we preset a main theorem, that is, let an inclusion $P \subset A$ of separable simple nuclear unital C*-algebras be of index finite type with the tracial Rokhlin property, and suppose that A is $\mathcal{Z}$-absorbing, then so does P (Theorem 5.4). As an application, any fixed point algebra A^H for any subgroup H of a finite group G is $\mathcal{Z}$-absorbing under the assumption that an action α from G on a simple nuclear unital C*-algebra A such that A is $\mathcal{Z}$-absorbing (Corollary 5.5). Before treating the strict comparison for a Cuntz semigroup, we consider the Cuntz equivalent for positive elements and show that under the assumption that an inclusion $P \subset A$ of unital C*-algebras has the tracial Rokhlin property, if for $n \in \mathbb{N}$ and positive

elements $a, b \in M_n(P)$ a is subequivalent to b in $M_n(A)$, then a is subequivalent to b in $M_n(P)$ (Proposition 6.2). Finally, we consider the strict comparison property for a Cuntz semigroup and show that the strict comparison property is inherited to P when an inclusion $P \subset A$ of simple separable exact unital C*-algebras has the tracial Rokhlin property and A has the strict comparison (Theore 7.2). Using the similar argument we show that if an inclusion $P \subset A$ of separable simple unital C*-algebras has the tracial Rokhlin property and the order on projections in A is determined by traces, then the order on projections in P is determined by traces (Corollary 7.3).

2. Local tracial $\mathcal{C}$-algebra

We recall the definition of local $\mathcal{C}$-property in [20].

Definition 2.1. Let $\mathcal{C}$ be a class of separable unital C*-algebras. Then $\mathcal{C}$ is *finitely saturated* if the following closure conditions hold:

(1) If $A \in \mathcal{C}$ and $B \cong A$, then $B \in \mathcal{C}$.
(2) If $A_1, A_2, \ldots, A_n \in \mathcal{C}$ then $\bigoplus_{k=1}^{n} A_k \in \mathcal{C}$.
(3) If $A \in \mathcal{C}$ and $n \in \mathbb{N}$, then $M_n(A) \in \mathcal{C}$.
(4) If $A \in \mathcal{C}$ and $p \in A$ is a nonzero projection, then $pAp \in \mathcal{C}$.

Moreover, the *finite saturation* of a class $\mathcal{C}$ is the smallest finitely saturated class which contains $\mathcal{C}$.

Definition 2.2. Let $\mathcal{C}$ be a class of separable unital C*-algebras. A *unital local $\mathcal{C}$-algebra* is a separable unital C*-algebra A such that for every finite set $S \subset A$ and every $\varepsilon > 0$, there is a C*-algebra B in the finite saturation of $\mathcal{C}$ and a unital *-homomorphism $\varphi\colon B \to A$ (not necessarily injective) such that $\mathrm{dist}(a, \varphi(B)) < \varepsilon$ for all $a \in S$.

When B in Definition 2.2 is non unital, we perturb the condition as follows:

Definition 2.3. Let $\mathcal{C}$ be a class of separable unital C*-algebras.

(i) A unital C*-algebra A is said to be a unital local tracial $\mathcal{C}$-algebra ([36]) if for every finite set $\mathcal{F} \subset A$ and every $\varepsilon > 0$, and any non-zero $a \in A^+$, there exist a non-zero projection $p \in A$, a sub C*-algebra $C \subset A$, and *-homomorphism $\varphi\colon C \to A$ such that $C \in \mathcal{C}$, $\varphi(1_C) = p$, and for all $x \in \mathcal{F}$,
   (1) $\|xp - px\| < \varepsilon$,
   (2) $pxp \in_\varepsilon \varphi(C)$, and
   (3) $1 - p$ is Murray-von Neumann equivalent to a projection in $\overline{aAa}$.

(ii) A unital C*-algebra A is said to belong to the class TA$\mathcal{C}$ ([4]) if for every finite set $\mathcal{F} \subset A$ and every $\varepsilon > 0$, and any non-zero $a \in A^+$, there exist a non-zero projection $p \in A$ and a sub C*-algebra $C \subset A$ such that $C \in \mathcal{C}$, $1_C = p$, and for all $x \in \mathcal{F}$,
   (1) $\|xp - px\| < \varepsilon$,
   (2) $pxp \in_\varepsilon C$, and
   (3) $1 - p$ is Murray-von Neumann equivalent to a projection in $\overline{aAa}$.

Remark 2.4. (1) When a unital C*-algebra A is a unital local tracial $\mathcal{C}$-algebra and each $\varphi(C) \in \mathcal{C}$, A belongs to the class TA$\mathcal{C}$.

(2) If $\mathcal{C}$ is the set of finite deimensional C*-algebras $\mathscr{F}$, and the class of interval algebras $\mathscr{I}$, respectively, then a local TA$\mathscr{F}$-algebra belongs to the class of C*-algebras of tracially AF C*-algebras ([14]), and a local TA$\mathscr{I}$-algebra belongs to the class of tracial topological one (TAI- algebras) ([16]) in the sense of Lin, respectively.

We have the following relation between the local $\mathcal{C}$ property and the local tracial approximational $\mathcal{C}$ property.

Recall that a C*-algebra A is said to have the *Property (SP)* if any nonzero hereditary C*-subalgebra of A has a nonzero projection.

Proposition 2.5. Let $\mathcal{C}$ be a finite saturated set and A be a local tracial $\mathcal{C}$-algebra. Then A has the property (SP) or A is a local $\mathcal{C}$ algebra.

Proof. Suppose that A does not have the Property (SP). Then there is a positive element $a \in A$ such that $\overline{aAa}$ has no non zero projection. Since A is a local TA$\mathcal{C}$-algebra, for every finite set $\mathcal{F} \subset A$ and every $\varepsilon > 0$, we conclude that there are a unital C*-algebra C in the class $\mathcal{C}$ and a unital *-homomorphism $\varphi\colon C \to A$ such that $\mathcal{F}$ can be approximated by a C*-algebra $\varphi(C)$ to whithin ε. Hence A is a local $\mathcal{C}$-algebra. ∎

3. Tracial Rokhlin property for finite group actions

Inspired by the concept of the tracial AF C*-algebras in [14] Phillips defined the tracial Rokhlin property for a finite group action in [24, Lemma 1.16] as follows:

Definition 3.1. Let α be the action of a finite group G on a unital an infinite dimensional finite simple separable unital C*-algebra A. An α is said to have the *tracial Rokhlin property* if for every finite set $F \subset A$, every $\varepsilon > 0$, and every nonzero positive $x \in A$, there are mutually orthogonal projections $e_g \in A$ for $g \in G$ such that:

(1) $\|\alpha_g(e_h) - e_{gh}\| < \varepsilon$ for all $g, h \in G$.
(2) $\|e_g a - a e_g\| < \varepsilon$ for all $g \in G$ and all $a \in F$.
(3) With $e = \sum_{g\in G} e_g$, the projection $1 - e$ is Murray-von Neumann equivalent to a projection in the hereditary subalgebra of A generated by x.

It is obvious that the Rokhlin property is stronger than the tracial Rokhlin property. As pointed in [9] the Rokhlin property gives rise to several K-theoretical constrains. For example there is no action with the Rokhlin property on the non-commutative 2-torus. On the contrary, if A is a simple higher dimensional non-commutative torus, which standard unitary generators $u_1, u_2, \ldots, u_n$, then the automorphism which sends u_k to $\exp(2\pi/n)u_k$, and fixes u_j for $j \neq k$, generates an action $\mathbb{Z}/n\mathbb{Z}$ and has the tracial Rokhlin property, but for $n > 1$ never has the Rokhlin property ([24]).

Lemma 3.2. ([24], [1]) Let a be an infinite dimensional stably finite simple C*-algebra with the Property SP. Let G be a finite group of order n and let $\alpha\colon G \to$

$\mathrm{Aut}(A)$ be an action of G with the tracial Rokhlin property. Then for any $\varepsilon > 0$, every $N \in \mathbb{N}$, any finite set $\mathcal{F} \subset A \rtimes_\alpha G$, and any non-zero $z \in (A \rtimes_\alpha G)^+$, there exist a non-zero projection $e \in A \subset A \rtimes_\alpha G$, a unital C*-subalgebra $D \subset e(A \rtimes_\alpha G)e$, a projection $f \in A$ and an isomorphism $\phi\colon M_n \otimes fAf \to D$, such that

- For every $a \in \mathcal{F}$ $\|ea - ae\| < n\varepsilon$.
- With (e_{gh}) for $g, h \in G$ being a system of matrix units for M_n, we have $\phi(e_{11} \otimes a) = a$ for all $a \in fAf$ and $\phi(e_{gg} \otimes 1) \in A$ for $g \in G$.
- With (e_{gg}) as in (1), we have $\|\phi(e_{gg} \otimes a) - \alpha_g(a)\| \leq \varepsilon\|a\|$ for all $a \in fAf$.
- For every $a \in F$ there exist $b_1, b_2 \in D$ such that $\|ea - b_1\| < \varepsilon$, $\|ae - b_2\| < \varepsilon$ and $\|b_1\|, \|b_2\| \leq \|a\|$.
- $e = \sum_{g \in G} \phi(e_{gg} \otimes 1)$.
- $1 - e$ is Murray-von Neumann equivalent to a projection in $\overline{z(A \rtimes_\alpha G)z}$.

Proof. This comes from the same argument in Lemma 3.1 of [1]. ∎

Theorem 3.3. Let $\mathcal{C}$ be a class of infinite dimensional stably finite separable unital C*-algebras which is closed under the following conditions:

(1) $A \in \mathcal{C}$ and $B \cong A$, then $B \in \mathcal{C}$.
(2) If $A \in \mathcal{C}$ and $n \in \mathbb{N}$, then $M_n(A) \in \mathcal{C}$.
(3) If $A \in \mathcal{C}$ and $p \in A$ is a nonzero projection, then $pAp \in \mathcal{C}$.

For any simple C*-algebra $A \in TA\mathcal{C}$ and an action α of a finite group G if α has the tracial Rokhlin property, then the crossed product algebra $A \rtimes_\alpha G$ belongs to the class $TA\mathcal{C}$.

Proof. Since α is outer by Lemma 1.5 of [24], $A \rtimes_\alpha G$ is simple by Theorem 3.1 of [12].

By Lemma 1.13 of [24], A has the Property (SP) or α has the strictly Rokhlin property.

If α has the strictly Rokhlin property, then $A \rtimes_\alpha G \in TA\mathcal{C}$ from the argument in the proof of Proposition 1.7 in [20].

Let $\mathcal{F} \subset A \rtimes_\alpha G$ be a finite set with $\|z\| \leq 1$ and let $\varepsilon > 0$.

Next, suppose that A has the property SP. Then there exists a non-zero projection $q \in A$ which is Murray-von Neumann equivalent in $A \rtimes_\alpha G$ to a projection in $\overline{z(A \rtimes_\alpha G)z}$ by Theorem 2.1 of [19]. Since A is simple, take orthogonal projections q_1, q_2 with $q_1, q_2 \leq q$ by [15]. Set $n = card(G)$, and set $\varepsilon_0 = \frac{1}{16n}\varepsilon$.

By Lemma 3.2 for n as given, for ε_0 in place of ε, and for q_1 in place of z there exist a non-zero projection $e \in A \subset A \rtimes_\alpha G$, a unital C*-subalgebra $D \subset e(A \rtimes_\alpha G)e$, a projection $f \in A$ and an isomorphism $\phi\colon M_n \otimes fAf \to D$, such that

(0) For every $a \in \mathcal{F}$ $\|ea - ae\| < n\varepsilon_0$.
(i) With (e_{gh}) for $g, h \in G$ being a system of matrix units for M_n, we have $\phi(e_{11} \otimes a) = a$ for all $a \in fAf$ and $\phi(e_{gg} \otimes 1) \in A$ for $g \in G$.
(ii) With (e_{gg}) as in (1), we have $\|\phi(e_{gg} \otimes a) - \alpha_g(a)\| \leq \varepsilon_0$ for all $a \in fAf$.
(iii) For every $a \in F$ there exist $d_1, d_2 \in D$ such that $\|ea - d_1\| < \varepsilon_0$, $\|ae - d_2\| < \varepsilon_0$ and $\|d_1\|, \|d_2\| \leq 1$.
(iv) $e = \sum_{g \in G} \phi(e_{gg} \otimes 1)$.
(v) $1 - e$ is Murray-von Neumann equivalent to a projection in $\overline{q_1(A \rtimes_\alpha G)q_1}$.

We note that there is a finite set T in the closed ball of $M_n \otimes fAf$ such that for every $a \in F$ there are $b_1, b_2 \in T$ such that $\|ea - \phi(b_1)\| < \varepsilon_0$ and $\|ae - \phi(b_2)\| < \varepsilon_0$.

Since A is simple, we choose equivalent nonzero projections $f_1, f_2 \in A$ such that $f_1 \preceq f$ and $f_2 \preceq q_2$ by [15]. Since $M_n \otimes fAf \in TA\mathcal{C}$ by Lemma 2.3 of [4], there is a projection $p_0 \in M_n \otimes fAf$ and a C*-subalgebra $C \subset M_n \otimes fAf$ such that $C \in \mathcal{C}$, $1_C = p_0$ such that $\|p_0 b - bp_0\| < \frac{1}{4}\varepsilon$ for all $b \in T$, such that for every $b \in T$ there exists $c \in C$ with $\|p_0 b p_0 - c\| < \frac{1}{4}\varepsilon$, and such that $1 - p_0 \preceq e_{11} \otimes f_1$ in $M_n \otimes fAf$.

Set $p = \phi(p_0)$, and set $E = \phi(C)$, which is a unital subalgebra of $p(A \rtimes_\alpha G)p$ and belongs to $\mathcal{C}$. Note that $e - p = \phi(1 - p_0) \preceq \phi(e_{11} \otimes f_1) = f_1$.

Let $a \in \mathcal{F}$. Choose $b \in T$ such that $\|\phi(b) - eae\| < \varepsilon_0 < \frac{1}{4}\varepsilon$. Then, using $pe = ep = p$,

$$\begin{aligned}\|pa - ap\| &\le 2\|ea - ae\| + \|peae - eaep\| \\ &\le 2\|ea - ae\| + 2\|eae - \phi(b)\| + \|p_0 b - bp_0\| \\ &< 2n\varepsilon_0 + 2\varepsilon_0 + \varepsilon_0 \\ &= (2n+3)\varepsilon_0 < \varepsilon.\end{aligned}$$

Choosing $c \in C$ such that $\|p_0 b p_0 - c\| < \frac{1}{4}\varepsilon$, the element $\phi(c)$ is in E and satisfies

$$\begin{aligned}\|pap - \phi(c)\| &= \|peaep - \phi(c)\| \\ &= \|p(eae - \phi(b))p + p\phi(b)p - \phi(c)\| \\ &\le \|eae - \phi(b)\| + \|p_0 b p_0 - c\| \\ &< \frac{1}{4}\varepsilon + \frac{1}{4}\varepsilon < \varepsilon.\end{aligned}$$

Finally, in $A \rtimes_\alpha G$ we have

$$1 - p = (1 - e) + (e - p) \preceq q_1 + q_2 \le q$$

and q is Murray-von Neumann equivalent to a projection in $\overline{z(A \rtimes_\alpha G)z}$. ∎

Theorem 3.4. ([6]) Let A be an infinite dimensional simple separable unital C*-algebra with stable rank one and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G with the tracial Rokhlin property. Then $A \rtimes_\alpha G$ has stable rank one.

Proof. Let $\mathcal{C}$ be the set of unital C*-algebras with stable rank one. Then $\mathcal{C}$ is closed under three conditions in Theorem 3.3. Then from Theorem 3.3 $A \rtimes_\alpha G$ belongs to the class TA$\mathcal{C}$.

Hence from Therem 4.3 of [4] $A \rtimes_\alpha G$ has stable rank one. ∎

We can also show that when $\mathcal{C}$ is the class of unital separable C*-algberas with real rank zero and A is a simple unital C*-algebra in the class TA$\mathcal{C}$, A has real rank zero.

Theorem 3.5. Let $\mathcal{C}$ be the class of unital separable C*-algberas with real rank zero. Then any simple unital C*-algebra in the class TA$\mathcal{C}$ has real rank zero.

Proof. We can deduce it from the same argument in the proof of Theorem 4.3 in [4]. ∎

Corollary 3.6. ([6]) Let A be an infinite dimensional simple separable unital C*-algebra with real rank zero and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G with the tracial Rokhlin property. Then $A \rtimes_\alpha G$ has real rank zero.

Proof. Let $\mathcal{C}$ be the set of unital C*-algebras with real rank zero. Then $\mathcal{C}$ is closed under three conditions in Theorem 3.3. Then from Theorem 3.3 $A \rtimes_\alpha G$ belongs to the class TA$\mathcal{C}$.

Hence from Theorem 3.5 $A \rtimes_\alpha G$ has real rank zero. ∎

Theorem 3.7. Let A be an infinite dimensional simple separable unital C*-algebra such that the order on projections over A is determined by traces and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G with tracial Rokhlin property. Then the order on projections over $A \rtimes_\alpha G$ is determined by traces.

Proof. Let $\mathcal{C}$ be the set of unital C*-algebras such that the order on projections over them is determined by traces. Then $\mathcal{C}$ is closed under three conditions in Theorem 3.3. Then from Theorem 3.3 $A \rtimes_\alpha G$ belongs to the class TA$\mathcal{C}$.

Hence from Therem 4.12 of [4] the order on projections over $A \rtimes_\alpha G$ is determined by traces. ∎

Definition 3.8. Let $\mathcal{T}^{(0)}$ be the calss of all finite dimensional C*-algebras and let $\mathcal{T}^{(k)}$ be the class of all C*-algebras with the form $pM_n(C(X))p$, where X is a finite CW complex with dimension k and $p \in M_n(C(X))$ is a projection.

A simple unital C*-algebra A is said to have tracial topological rank no more than k if for any set $\mathcal{F} \subset A$, and $\varepsilon > 0$ and any nonzero positive element $a \in A$, there exists a C*-subalgebra $B \subset A$ with $B \in \mathcal{T}^{(k)}$ and $id_B = p$ such that

(1) $\|xp - px\| < \varepsilon$

(2) $pxp \in_\varepsilon C$, and

(3) $1 - p$ is Murray-von Neumann equivalent to a projection in $\overline{aAa}$.

Theorem 3.9. ([21]) Let A be an infinite dimensional simple unital C*-algebra with tracial topological rank no more than or equal to k, and $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G with tracial Rokhlin property. Then $A \rtimes_\alpha G$ has tracial topological rank more than or equal to k.

Proof. Let $\mathcal{C}$ be the set $\mathcal{T}^{(k)}$. Then $\mathcal{T}^{(k)}$ is closed under three conditions in Theorem 3.3. Then from Theorem 3.3 $A \rtimes_\alpha G$ belongs to the class TA$\mathcal{C}$. This means that $A \rtimes_\alpha G$ has tracial topological rank no more that or equal to k from the Definition 3.8. ∎

4. Tracial Rokhlin property for a inclusion of unital C*-algebras

Let $P \subset A$ be a inclusion of unital C*-algebras and $E\colon A \to P$ be a conditional expectation of index finite.

As in the case of the Rokhlin property in [13] we can define the tracial Rokhlin property for a conditional expectation for an inclusion of unital C*-algebras.

Definition 4.1. Let $P \subset A$ be an inclusion of unital C*-algebras and $E\colon A \to P$ be a conditional expectation of index finite. A conditional expectation E is said to have the *tracial Rokhlin property* if for any nonzero positive $z \in A^\infty$ there exists a projection $e \in A' \cap A^\infty$ satisfying

$$(\mathrm{Index}E)E^\infty(e) = g$$

is a projection and $1 - g$ is Murray-von Neumann equivalent to a projection in the hereditary subalgebra of A^∞ generated by z, and a map $A \ni x \mapsto xe$ is injective. We call e a Rokhlin projection.

As in the case of an action with the tracial Roklin property ([24, Lemma 1.13]) if $E\colon A \to P$ is a conditional expectation of index finite type for an inclusion of unital C*-algebras $P \subset A$ and E has the tracial Rokhlin property, then A has Property (SP) or E has the Rokhlin property, that is, there is Rokhlin projection $e \in A' \cap A^\infty$ such that $(\mathrm{Index}E)E^\infty(e) = 1$.

Lemma 4.2. Let $P \subset A$ be an inclusion of unital C*-algebras and $E\colon A \to P$ be a conditional expectation of index finite type. Suppose that E has the tracial Rokhlin property, then A has the Property (SP) or E has the Rokhlin property.

Proof. If A does not have the Property (SP), then A^∞ does not have Property (SP), that is, there is a nonzero positive element $x \in A^\infty$ which generates a hereditary subalgebra which contains no nonzero projection. Since E has the tracial Rokhlin property, there exists a projection $e \in A_\infty$ such that $1 - (\mathrm{Index}E)E^\infty(e)$ is equivalent to some projection in $\overline{xA^\infty x}$. Hence $1 - (\mathrm{Index}E)E^\infty(e) = 0$. This implies that E has the Rokhlin property. ∎

Remark 4.3. (1) When A has the Property (SP), we know that g in Definition 4.1 is not zero. Indeed, since A^∞ has the Property (SP), from [24, Lemma 1.15] for a positive element $x \in A^\infty$ with $\|x\| = 1$ and $\varepsilon > 0$ there exists a nonzero projection $q \in \overline{xA^\infty x}$ such that, whenever $1 - g \preceq q \in \overline{xA^\infty x}$, then

$$\|gxg\| > 1 - \varepsilon.$$

This implies that $g \neq 0$.

(2) We always have $g \in P' \cap P^\infty$. Indeed, for any $x \in P$

$$\begin{aligned} xg &= x(\mathrm{Index}E)E^\infty(e) \\ &= (\mathrm{Index}E)E^\infty(xe) \\ &= (\mathrm{Index}E)E^\infty(ex) \\ &= (\mathrm{Index}E)E^\infty(e)x \\ &= gx. \end{aligned}$$

Remark 4.4. In Definition 4.1 when A is simple,

(1) we do not need the injectivity of the map $A \ni x \mapsto xe$.

(2) we have $ege = e$. Indeed, since A is simple, Index$E$ is scalar by [33, Remark 2.3.6] and from [33, Lemma 2.1.5 (2)] we have

$$E(e) \geq \frac{1}{(\mathrm{Index}E)^2}e,$$

hence $g \geq \frac{1}{\mathrm{Index}E}e$. Then

$$\begin{aligned}(1-g)\frac{1}{\mathrm{Index}E}e(1-g) &= 0\\ (1-g)e &= 0\\ e &= ge = ege\end{aligned}$$

(3) Let $E\colon A \to P$ be of index finite with the tracial Rokhlin property and let consider the basic extension

$$P \subset A \subset B.$$

Then the Rokhlin projection $e \in A' \cap A^\infty$ satisfies $eBe = Ae$.

Indeed, let e_p be the Jones projection for the inclusion $A \supset P$. Set $f = \mathrm{Index}Eee_pe$. Then

$$\begin{aligned}f^2 &= (\mathrm{Index}E)^2 ee_pee_pe\\ &= (\mathrm{Index}E)^2 eE^\infty(e)e_pe\\ &= (\mathrm{Index}E)^2 eE^\infty(e)e_pe\\ &= (\mathrm{Index}E)ege_pe\\ &= (\mathrm{Index}E)ee_pe\\ &= f.\end{aligned}$$

Since

$$\begin{aligned}\hat{E}^\infty(e-f) &= e - \mathrm{Index}E\hat{E}^\infty(ee_pe)\\ &= e - e = 0,\end{aligned}$$

we have $e = f = \mathrm{Index}Eee_pe$, $\hat{E}$ is the dual conditional expectation for E.

Then we have for any $x, y \in A$

$$\begin{aligned}e(xe_py)e &= xee_pey\\ &= (\mathrm{Index}E)^{-1}xey\\ &= (\mathrm{Index}E)^{-1}xye \in Ae.\end{aligned}$$

Hence $eBe \subset Ae$. Conversely, since $A \subset B$, $Ae \subset eBe$, and we conclude that $eBe = Ae$. ∎

The following is the heredity of the Property SP for an inclusion of unital C*-algebras.

Proposition 4.5. Let $P \subset A$ be an inclusion of unital C*-algebras with index finite type. Suppose that A is simple and $E\colon A \to P$ has the tracial Rokhlin property. Then we have

(1) P is simple.
(2) A has the Property (SP) if and only if P has the Property (SP).

Proof. (1): Let e be a Rokhlin projection for E and $P \subset A \subset B$ be the basic extension. Since P is stably isomorphic to B, we will show that B is simple, that is, $B' \cap B = \mathbb{C}$.

Since $e = [(e_n)] \in A' \cap A^\infty$, for any $x \in A' \cap B$ we have

$$ex = [(e_n)]x = [(e_n x)] = [(xe_n)] = xe.$$

We may assume that $x = a_1 e_p a_2$, where e_p is the Jones projection for E. Then

$$\begin{aligned} xe &= exe \\ &= e(a_1 e_p a_2)e \\ &= a_1 e e_p e a_2 \\ &= (\mathrm{Index}E)^{-1} a_1 a_2 e \\ &= \hat{E}(x)e, \end{aligned}$$

where $\hat{E}\colon B \to A$ be the dual conditional expectation of E. Note that $\hat{E}(x) \in A'$. Hence we have $xe \in (A' \cap A)e$.

Since A is simple and $(B' \cap B)e \subset (A' \cap B)e \subset (A' \cap A)e$, we have $(B' \cap B)e = \mathbb{C}e$. Since the map $\rho\colon A' \cap B \to (A' \cap B)e$ by $\rho(x) = xe$ is an isomorphism, $B' \cap B = \mathbb{C}$, that is, B is simple, and P is simple.

(2): It follows from [19]. ∎

Proposition 4.6. Let G be a finite group, α an action of G on an infinite dimensional finite simple separable unital C*-algebra A, and E the canonical conditional expectation from A onto the fixed point algebra $P = A^\alpha$ defined by

$$E(x) = \frac{1}{|G|} \sum_{g \in G} \alpha_g(x) \quad \text{for } x \in A,$$

where $|G|$ is the order of G. Then α has the tracial Rokhlin property if and only if E has the tracial Rokhlin property.

Proof. Suppose that α has the tracial Rokhlin property. Since A is separable, there is an increasing sequence of finite sets $\{F_n\}_{n \in \mathbb{N}} \subset A$ such that $\overline{\cup_{n \in \mathbb{N}} F_n} = A$. Let any nonzero positive element $x = (x_n) \in A^\infty$. Then we may assume that each x_n is nonzero positive element. The simplicity of A implies that the map $A \ni x \mapsto xe$ is injective.

Since α has the tracial Rokhlin property, for each n there are mutually orthogonal projections $\{e_{g,n}\}_{g \in G}$ such that

$$\|\alpha_h(e_{g,n}) - e_{hg,n}\| < \frac{1}{n}$$

for all $g, h \in G$,

$$\|[e_{g,n}, a]\| < \frac{1}{n}$$

for all $g \in G$, and $a \in F_n$ with $\|a\| \leq 1$

$$1 - \sum_{g \in G} e_{g,n}$$

is equivalent to a nonzero projection q_n in $\overline{x_n A x_n}$.

Set $e_g = [(e_{g,n})] \in A^\infty$ for $g \in G$. Then for all $g, h \in G$

$$\|\alpha_h^\infty(e_g) - e_{hg}\| = \limsup \|\alpha_h(e_{g,n}) - e_{hg,n}\| = 0,$$

hence $\alpha_h^\infty(e_g) = e_{hg}$ for all $g, h \in G$.

For all $a \in \cup_{n\in\mathbb{N}} F_n$ with $\|a\| \leq 1$ and all $g \in G$ we have

$$\|[e_g, a]\| = \limsup \|[e_{g,n}, a]\| = 0,$$

hence $e_g \in A_\infty$ for all $g \in G$.

Set $q = (q_n) \in A^\infty$. Then q is nonzero projection in $\overline{xA^\infty x}$ and

$$1 - \sum_{g\in G} e_g = (1 - \sum_{g\in G} e_{g,n}) \sim (q_n) = q.$$

Therefore, if we set $e = e_1$ for the identity element 1 in G, then $e \in A' \cap A^\infty$ and

$$\begin{aligned} E^\infty(e) &= \frac{1}{|G|} \sum_{g\in G} \alpha_g^\infty(e) \\ &= \frac{1}{|G|} \sum_{g\in G} e_g \end{aligned}$$

and

$$1 - |G|E^\infty(e) = 1 - \sum_{g\in G} e_g \sim q \in \overline{xA^\infty x}.$$

It follows that E has the tracial Rokhlin property.

Conversely, suppose that E has the tracial Rokhlin property. From Lemma 4.2 A has the Propety (SP) or E has the Rokhlin property. If E has the Rokhlin property, then α has the Rokhlin property by Proposition 3.2 in [13], hence α has the tracial Rokhlin property from the definition.

We may assume that A has Property (SP). Then for any finite set $F \subset A$, $\varepsilon > 0$, and any nonzero positive element $x \in A$ there is a projection $e \in A_\infty$ such that $|G|E^\infty(e)(= g)$ is a projection and $1-g$ is equivalent to a projection $q \in \overline{xA^\infty x}$. We note that $g \neq 0$ by Remark 4.3, and $e \neq 0$. When we write $e = (e_n)$ and $q = (q_n)$, we may assume that for each $n \in \mathbb{N}$ e_n is projection and $1 - e_n$ is equivalent to q_n.

Define $e_g = \alpha_g^\infty(e) \in A_\infty$ for $g \in G$, write $e_g = [(\alpha_g(e_n))] = [(e_{g,n})]$ for $g \in G$. Then since we have

$$\sum_{g\in G} e_g = \sum_{g\in G} \alpha_g(e) = |G|E^\infty(e) = g$$

and g is projection, we may assume that that $\{e_{g,n}\}_{g\in G}$ are mutually orthogonal projections for each $n \in \mathbb{N}$ by [15, Lemma 2.5.6].

Then $\alpha_h^\infty(e_g) = e_{hg}$ for all $g, h \in G$, $\|[e_g, a]\| = 0$ for all $a \in F$ and all $g \in G$, and

$$\begin{aligned} 1 - \sum_{g\in G} e_g &= 1 - \sum_{g\in G} \alpha_g^\infty(e) \\ &= 1 - |G|E(e) \\ &= 1 - g \sim q \in \overline{xA^\infty x}, \end{aligned}$$

Then there exists $n \in \mathbb{N}$ such that

$$\|\alpha_h(e_{g,n}) - e_{hg,n}\| < \varepsilon$$

for all $g, h \in G$,

$$\|[e_{g,n}, a]\| < \varepsilon$$

for all $a \in F$ and $g \in G$, and

$$1 - \sum_{g \in G} e_{g,n} \sim q_n \in \overline{xAx}.$$

Set $f_g = e_{g,n}$ for $g \in G$, then we have

$$\|\alpha_h(f_g) - f_{hg}\| < \varepsilon,$$

for all $g, h \in G$,

$$\|[f_g, a]\| < \varepsilon$$

for all $a \in F$ and $g \in G$, and

$$1 - \sum_{g \in G} f_g \sim q_n \in \overline{xAx}.$$

Hence α has the tracial Rokhlin property. ∎

The following is a key lemma to prove the main theorem in this section.

Lemma 4.7. Let $A \supset P$ be an inclusion of unital C*-algebras and E a conditional expectation from A onto P with index finite type. Suppose that A is simple. If E has the tracial Rokhlin property with a Rokhlin projection $e \in A_\infty$ and a projection $g = (\mathrm{Index}E)E^\infty(e)$, then there is a unital linear map $\beta\colon A^\infty \to P^\infty g$ such that for any $x \in A^\infty$ there exists the unique element y of P^∞ such that $xe = ye = \beta(x)e$ and $\beta(A' \cap A^\infty) \subset P' \cap P^\infty g$. In particular, $\beta_{|_A}$ is a unital injective *-homomorphism and $\beta(x) = xg$ for all $x \in P$.

Proof. Since E has the tracial Rokhlin property, A has the Property (SP) or E has the Rokhlin property by Lemma 4.2. If E has the Rokhlin property, then the conclusion comes from Lemma 2.5 in [22] with $g = 1$. Therefore we may assume that A has the Property (SP).

Since A has the Property (SP), g and e are nonzero projections by Remark 4.3. As in the same argument in the proof of Lemma 2.5 in [22] we have for any element x in A^∞ there exists a unique element $y = (\mathrm{Index}E)E^\infty(xe) \in P^\infty$ such that $xe = ye$. Note that since $eg = e$ by Remark 4.4, we have

$$\begin{aligned} yg &= (\mathrm{Index}E)E^\infty(xe)g \\ &= (\mathrm{Index}E)E^\infty(xeg) \\ &= (\mathrm{Index}E)E^\infty(xe) = y. \end{aligned}$$

Then we can define a unital map $\beta\colon A^\infty \to P^\infty g$ such that $xe = ye = \beta(x)e$ and $\beta(A' \cap A^\infty) \subset P' \cap P^\infty g$.

Note that β is injective. Indeed, if $\beta(x) = 0$ for $x \in A$ xe $= 0$. Hence from the definition of the tracial Rokhlin property for E, $x = 0$.

Since for any $x \in A$

$$\begin{aligned}\beta(x)g &= (\mathrm{Index}E)E^{\infty}(xe)g \\ &= (\mathrm{Index}E)E^{\infty}(xeg) \\ &= (\mathrm{Index}E)E^{\infty}(ex)\ (=\beta(x)) \\ &= (\mathrm{Index}E)E^{\infty}(gex) \\ &= g(\mathrm{Index}E)E^{\infty}(xe) \\ &= g\beta(x),\end{aligned}$$

we know that $\beta_{|A}$ is a unital *-homomorphism from A to $gP^{\infty}g$ from the same argument as in the proof of Lemma 2.5 in [22]. In particular for any $x \in P$ we have

$$\begin{aligned}\beta(x) &= (\mathrm{Index}E)E^{\infty}(xe) \\ &= x(\mathrm{Index}E)E^{\infty}(e) \\ &= xg(=gx).\end{aligned}$$

∎

Proposition 4.8. Let $P \subset A$ be an inclusion of unital C*-algebras with index finite type and $E\colon A \to P$ has the tracial Rokhlin property. Suppose that A is simple with $\mathrm{tsr}(A) = 1$. Then $\mathrm{tsr}(P) = 1$.

Proof. Since E has the tracial Rokhlin property, E has the Rokhlin property or A has the Property (SP) by Lemma 4.2. If E has the Rokhlin property, we conclude that $\mathrm{tsr}(P) = 1$ by [13]. Therefore, we assume that A has the Property (SP). Then we know that P is simple and has the Property (SP) by Proposition 4.5.

Since A is stably finite and an inclusion $P \subset A$ is of index-finite type, P is stably finite. Hence, using the idea in [25] we have only to show that any two sided zero divisor in P is approximated by invertible elements in P.

Let $x \in P$ be a two sided zero diviser. From [25, Lemma 3.5] we may assume that there is a positive element $y \in P$ such that $yx = 0 = xy$. Since P has the Property (SP), there is a non-zero projection $e \in \overline{yPy}$. Since P is simple, we can take orthogonal projections e_1 and e_2 in P such that $e = e_1 + e_2$ and $e_2 \preceq e_1$. Note that $x \in (1-e_1)A(1-e_1)$. Since $\mathrm{tsr}((1-e_1)A(1-e_1)) = 1$, there is an invertible element b in $(1-e_1)A(1-e_1)$ such that $||x-b|| < \dfrac{1}{3}\varepsilon$.

Since E has the tracial Rokhlin property, there is a projection $g \in P' \cap P^{\infty}$ such that $1-g \preceq e_2$. That is, there is a partial isometry $w \in P^{\infty}$ such that $w^*w = 1-g$ and $ww^* \leq e_1$. Moreover, $||\beta(x) - \beta(b)|| < \dfrac{1}{3}\varepsilon$ by Lemma 4.7. Note that $\beta(b)$ is invertible in $gP^{\infty}g$.

Set

$$z = \frac{\varepsilon}{3}(e_1 - ww^*) + \frac{\varepsilon}{3}w + \frac{\varepsilon}{3}w^* + (1-g)x(1-g).$$

Then z is invertible in $e_1P^{\infty}e_1 + (1-g)(1-e_1)P^{\infty}(1-e_1)(1-g)$ and $||z - (1-g)x(1-g)|| < \dfrac{\varepsilon}{3}$.

Then $\beta(b)+z \in P^\infty$ is invertible and

$$\begin{aligned} ||x-(\beta(b)+z)|| &= ||xg+x(1-g)-\beta(b)-z|| \\ &= ||\beta(x)-\beta(b)+(1-g)x(1-g)-z|| \\ &\le ||\beta(x)-\beta(b)|| + ||(1-g)x(1-g)-z|| \\ &\le \frac{\varepsilon}{3}+\frac{\varepsilon}{3} < \varepsilon. \end{aligned}$$

Write $\beta(b)+z=(y_n)$ such that y_n is invertible in P. Therefore, there is a y_n such that $||x-y_n|| < \varepsilon$, and we conclude that $\mathrm{tsr}(P)=1$. ∎

Proposition 4.9. Let $P \subset A$ be an inclusion of unital C*-algebras with index finite type and $E\colon A \to P$ has the tracial Rokhlin property. Suppose that A is simple with real rank zero. Then P has real rank zero.

Proof. Let $x \in P$ be a self-adjoint element and $\varepsilon > 0$. Consider a continuous real valued function f is defined by $f(x)=1$ for $|x| \le \frac{\varepsilon}{12}$, $f(t)=0$ if $|x| \ge \frac{\varepsilon}{64}$, and $f(t)$ is linear if $\frac{\varepsilon}{12} \le |x| \le \frac{\varepsilon}{6}$. We may assume that $f(x) \neq 0$. Note that $\|yx\| < \frac{\varepsilon}{6}$ for any $y \in \overline{f(x)Pf(x)}$.

Since A is simple and has the Property (SP), P has the Property (SP), that is, there is a non-zero projection $e \in \overline{f(x)Pf(x)}$. Moreover, there are orthogonal projections e_1 and e_2 such that $e=e_1+e_2$ such that $e_2 \preceq e_1$. Then

$$\begin{aligned} ||x-(1-e_1)x(1-e_1)|| &= ||e_1xe_1+e_1x(1-e_1)+(1-e_1)xe_1)|| \\ &< \frac{3\varepsilon}{12} = \frac{\varepsilon}{4} \end{aligned}$$

As in the same step in the argument in Proposition 4.8 we have there is an invertible self-adjoint element $z \in P$ such that $||(1-e_1)x(1-e_1)-z|| < \frac{2\varepsilon}{3}$. Hence, we have $||x-z|| < \varepsilon$, and we conclude that P has real rank zero. ∎

The following lemma is important to prove that heredity of the local tracial $\mathcal{C}$-property for an inclusion of unital C*-algerbas.

Lemma 4.10. Let $P \subset A$ be an inclusion of unital C*-algebras with index finite type and $E\colon A \to P$ has the tracial Rokhlin property. Suppose that projections $p, q \in P^\infty$ satisfy $ep=pe$ and $q \preceq ep$ in A^∞, where e is the Rokhlin projection for E. Then $q \preceq p$ in P^∞.

Proof. Let s be a partial isometry in A^∞ such that $s^*s=q$ and $ss^* \le ep$.

Set $v = (\mathrm{Index}E)^{1/2}E^\infty(s)$. Then

$$\begin{aligned} v^*ve_p &= \mathrm{Index}EE^\infty(s)^*E^\infty(s)e_p \\ &= \mathrm{Index}EE^\infty(s^*e)E^\infty(es)e_p \\ &= \mathrm{Index}Ee_ps^*ee_pese_p \\ &= e_ps^*ese_p \quad (\mathrm{Index}Eee_pe = e) \\ &= E^\infty(s^*es)e_p \\ &= E^\infty(s^*s)e_p \\ &= E^\infty(q)e_p \\ &= qe_p. \end{aligned}$$

Hence

$$\begin{aligned} \hat{E}^\infty(v^*ve_p) &= \hat{E}^\infty(qe_p) \\ (\mathrm{Index}E)^{-1}v^*v &= (\mathrm{Index}E)^{-1}q \\ v^*v &= q. \end{aligned}$$

Since

$$\begin{aligned} pv &= p(\mathrm{Index}E)^{1/2}E^\infty(s) \\ &= (\mathrm{Index}E)^{1/2}E^\infty(ps) \\ &= (\mathrm{Index}E)^{1/2}E^\infty(s) \\ &= v, \end{aligned}$$

we have $q \preceq p$ in P^∞. ∎

Theorem 4.11. Let $\mathcal{C}$ be a class of weakly semiprojective C*-algebras satisfying conditions in Theorem 3.3. Let $A \supset P$ be an inclusion of unital C*-algebras and E a conditional expectation from A onto P with index finite type. Suppose that A is simple, local tracial $\mathcal{C}$-algebra and E has the tracial Rokhlin property. Then P is a local tracial $\mathcal{C}$-algebra.

Proof. We shall prove that for every finite set $F \subset P$, every $\varepsilon > 0$, and $z \in P^+\backslash 0$ there are C*-algebra $Q \in \mathcal{C}$ with $q = 1_Q$ and *-homomorphism $\pi\colon Q \to A$ such that $\|\pi(q)x - x\pi(q)\| < \varepsilon$ for all $x \in F$, $\pi(q)S\pi(q) \subset_\varepsilon \pi(Q)$, and $1 - \pi(q)$ is a equivalent to some non-zero projection in $\overline{zPz}$.

Since E has the tracial Rokhlin property, A has the Property (SP) or E has the Rokhlin property by Lemma 4.2.

Suppose that E has the Rokhlin property. We have then from Lemma 2.5 in [22] there is a unital *-homomorphism $\beta\colon A \to P^\infty$ such that $\beta(x) = x$ for all $x \in P$. Since A is a local tracial $\mathcal{C}$-algebra, there is an algebra $B \in \mathcal{C}$ with $1_B = p$ and *-homomorphism $\pi\colon B \to A$ such that $\|x\pi(p) - \pi(p)x\| < \varepsilon$ for all $x \in F$, $\pi(p)F\pi(p) \subset_\varepsilon \pi(B)$, and $1 - \pi(p)$ is equivalent to a non-zero projection $q \in \overline{zAz}$. Since E has the Rokhlin property, there exists a non-zero projection $e \in A' \cap A^\infty$ such that $E^\infty(e) = \frac{1}{\mathrm{IndexE}}$.

Since B is weakly semiprojective, there exists $k \in \mathbb{N}$ and $\overline{\beta \circ \pi}\colon B \to \prod_{n=k}^\infty P$ such that $\beta \circ \pi = \pi_k \circ \overline{\beta \circ \pi}$, where $\pi_k((b_k, b_{k+1}, \dots)) = (0, \dots, 0, b_k, b_{k+1}, \dots)$. For each $l \in \mathbb{N}$ with $l \geq k$ let β_l a *-homomorphism from B to P so that $\overline{\beta \circ \pi}(b) =$

$(\beta_n(b))_{n=k}^{\infty}$ for $b \in B$. Then $\beta \circ \pi(b) = (0, \dots, 0, \beta_k(b), \beta_{k+1}(b), \dots) + C_0(P)$ for $b \in B$ and β_l is a *-homomorphism for $l \geq k$.

Since $1 - p \sim q \in \overline{zAz}$,

$$\begin{aligned} 1 - \beta \circ \pi(p) &= 1 - \beta(1 - \pi(p)) \\ &= \beta(1 - \pi(p)) \sim \beta \circ \pi(q) \in \beta(\overline{zAz}) \\ &[(1 - \beta_k(\pi(p)))] \sim [(q_k)] \in \overline{zP^{\infty}z}, \end{aligned}$$

where each q_k are projections in P. Taking the sufficient large k since $\lim_k \|\beta_k(x) - x\| = 0$ for $x \in P$, we have

(1) $\|x\beta_k(p) - \beta_k(p)x\| < 2\varepsilon$ for any $x \in F$,
(2) $\beta_k(p)F\beta_k(p) \subset_{\varepsilon} \beta_k(p)\beta_k(B)\beta_k(p)$, and
(3) $1 - \beta_k(p) = \beta_k(1 - p) \sim q_k \in \overline{zPz}$.

Hence P is a local tracially $\mathcal{C}$-algebra.

Suppose that A has the Property (SP). Since A is simple, from Proposition 4.5 P has also the Property (SP). Let $F \subset P$ be a finite set, $\varepsilon > 0$, and $z \in P^+ \backslash 0$. Since P is simple and has the Property (SP), there is orthogonal non-zero projections $r_1, r_2 \in \overline{zPz}$.

Since A is a local tracial $\mathcal{C}$-algebra, there is an algebra $B \in \mathcal{C}$ with $1_B = p$ and *-homomorphism $\pi \colon B \to A$ such that $\|x\pi(p) - \pi(p)x\| < \varepsilon$ for all $x \in F$, $\pi(p)F\pi(p) \subset_{\varepsilon} B$, and $1 - \pi(p)$ is equivalent to a non-zero projection $q \in \overline{r_1Ar_1}$. Since E has the tracial Rokhlin property, there exist the Rokhlin propjection $e' \in A' \cap A^{\infty}$. Take another Rokhlin projection $e \in A' \cap A^{\infty}$ for a projection $e'r_2$ such that $g = \mathrm{Index}EE(e)$ satifies $1 - g$ is equivalent to a projection $\overline{e'r_2A^{\infty}e'r_2}$, that is, $1 - g \preceq e'r_2$ in A^{∞}. By Lemma 4.10 we know, then, that $1 - g \preceq r_2$ in P^{∞}, that is, there is a projection $s \leq r_2 \in P^{\infty}$ such that $1 - g \sim s$.

Write $g = [(g_n)]$ for some projections $\{g_k\}_{k \in \mathbb{N}} \subset P$. From Lemma 4.7 there exists injective *-homomorphism $\beta \colon A \to gP^{\infty}g$ such that $\beta(x) = xg$ for all $x \in P$. Since B is weakly semiprojective, there exists $k \in \mathbb{N}$ and $\overline{\beta \circ \pi} \colon B \to \prod_{n=k}^{\infty} P$ such that $\beta \circ \pi = \pi_k \circ \overline{\beta \circ \pi}$, where

$$\pi_k((b_k, b_k + 1, \dots)) = (0, \dots, 0, b_k, b_{k+1}, \dots) + C_0(P).$$

For each $l \in \mathbb{N}$ with $l \geq k$ let β_l be a map from B to g_lPg_l so that $\overline{\beta \circ \pi}(b) = (\beta_l(b))_{l=k}^{\infty}$ for $b \in B$. Then $\beta \circ \pi(b) = (0, \dots, 0, \beta_k(b), \beta_{k+1}(b), \dots) + C_0(P)$ for all $b \in B$ and β_l is a *-homomorphism for $l \geq k$.

Since $1 - \pi(p) \sim q \in \overline{r_1Ar_1}$,

$$\begin{aligned} 1 - (\beta \circ \pi)(p) &= 1 - g + g - \beta(\pi(p)) \\ &= 1 - g + \beta(1 - \pi(p)) \\ &\sim s + \beta(q) \in r_2P^{\infty}r_2 + \iota \circ \beta(\overline{r_1Ar_1}) \\ &\subset r_2P^{\infty}r_2 + r_1P^{\infty}r_1 \subset \overline{zP^{\infty}z}, \end{aligned}$$

we have $[(1 - \beta_k(p)))] \sim [(q_k)] \in \overline{zP^{\infty}z}$,
where each q_k is projection in P. Taking the sufficient large k since $\lim_k \|\beta_k(x) - x\| = 0$ for $x \in P$, we have

(1) $\|x\beta_k(p) - \beta_k(p)x\| < 2\varepsilon$ for any $x \in F$,
(2) $\beta_k(p)F\beta_k(p) \subset_\varepsilon \beta_k(p)\beta_k(B)\beta_k(p)$, and
(3) $1 - \beta_k(p) \sim q_k \in \overline{zPz}$.

Hence P is a local tracially $\mathcal{C}$-algebra. ∎

Corollary 4.12. Let $P \subset A$ be an inclusion of unital C*-algebras and E a conditional expectation from A onto P with index finite type. Suppose that A is an infinite dimensional simple C*-algebra with tracial topological rank zero (resp. less than or equal to one) and E has the tracial Rokhlin property. Then P has tracial rank zero (resp. less than or equal to one).

Proof. Since the classes $\mathcal{T}^{(k)}$ $(k = 0, 1)$ are semiprojective with respect to a class of unital C*-algebras ([17]) and finitely saturated ([20, Examples 2.1 and 2.2, and Lemma 1.6]), the conclusion comes from Theorem 4.11 and Definition 3.8. ∎

5. Jiang-Su absorption

In this section we discuss about the heredity for the Jiang-Su absorption for an inclusion of unital C*-algebras with the tracial Rokhlin property.

Definition 5.1. ([8]) A unial C*-algebra A is said to be tracially $\mathcal{Z}$-absorbing if $A \ncong \mathbb{C}$ and for any finite set $F \subset A$ and non-zero positive element $a \in A$ and $n \in \mathbb{N}$ there is an order zero construction $\phi\colon M_n \to A$ such that the following hold:

(1) $1 - \phi(1) \preceq a$,
(2) For any normalized element $x \in M_n$ and any $y \in F$ we have $\|[\phi(x), y)]\| < \varepsilon$.

Theorem 5.2. ([8, Theorem 4.1]) Let A be a unital, separable, simple, nuclear C*-algebra. If A is tracially $\mathcal{Z}$-absorbing, then $A \cong A \otimes \mathcal{Z}$.

Note that for a simple unital C*-algebra A if A is $\mathcal{Z}$-absorbing, then A is tracially $\mathcal{Z}$-absorbing ([8, Proposition 2.2]).

Theorem 5.3. Let $P \subset A$ be an inclusion of unital C*-algebra and E be a conditional expectation from A onto P with index finite type. Suppose that A is simple, separable, unital, tracially $\mathcal{Z}$-absorbing and E has the tracial Rohklin property. Then P is tracially $\mathcal{Z}$-absorbing.

Proof. Take any finite set $F \subset P$ and non-zero positive element $a \in P$ and $n \in \mathbb{N}$. Since $E\colon A \to P$ has the tracial Rokhlin property, E has the Rokhlin property or A has the Property (SP). If E has the Rokhlin property, then P is $\mathcal{Z}$-absorbing ([22]), and we are done. Hence we may assume that A has the Property (SP).

Since A is simple and has the Property (SP), P has the Property (SP) by Proposition 4.5. Then there exists orthogonal projections p_1, p_2 in $\overline{aPa}$.

Since A is tracially $\mathcal{Z}$-absorbing, there is an order zero construction $\phi\colon M_n \to A$ such that the following hold:

(1) $1 - \phi(1) \preceq p_1$,
(2) For any normalized element $x \in M_n$ and any $y \in F$ we have $\|[\phi(x), y)]\| < \varepsilon$.

Since $E\colon A \to P$ has the tracial Rokhlin, there is a projection $e \in A' \cap A^\infty$ satisfying $(\mathrm{Index}E)E^\infty(e) = g$ is a projection and $1 - g \preceq p_2$. Moreover, there is an

injective *homomorphism β from A into $gP^{\infty}g$ such that $\beta(1) = g$ and $\beta(a) = ag$ for $a \in P$.

Then a function $\beta \circ \phi (= \psi)\colon M_n \to P^{\infty}$ is an order zero map such that

(i)
$$\begin{aligned} 1 - \psi(1) &= 1 - (\beta \circ \phi)(1) \\ &= 1 - g + \beta(1 - \phi(1)) \\ &\preceq p_2 + \beta(p_1) \\ &= p_2 + p_1\beta(1) \\ &\preceq a \end{aligned}$$
, that is, $1 - \psi(1) \preceq a$ in P^{∞}.

(ii) For any normalized element $x \in M_n$ and $y \in F$
$$\begin{aligned} \|[\psi(x), y]\| &= \|[\beta(\phi(x)), y]\| \\ &= \|\beta(\phi(x))y - y\beta(\phi(x))\| \\ &= \|\beta(\phi(x))\beta(y) - \beta(y)\beta(\phi(x))\| \\ &= \|\beta(\phi(x)y - y\phi(x))\| \\ &\leq \|\phi(x)y - y\phi(x)\| < \varepsilon. \end{aligned}$$

Since $C^*(\phi(M_n))$ is semiprojective by [34, Proposition 3.2 (a)], there is a $k \in \mathbb{N}$ and a *-homomorphism $\tilde{\beta}\colon C^*(\phi(M_n))) \to \Pi P/ \oplus_{i=1}^{k} P \to P^{\infty}$ such that $\pi_k \circ \tilde{\beta} = \beta$, where π_k be the canonical map from $\Pi P/ \oplus_{i=1}^{k} P$ to P^{∞}. Write $\tilde{\beta}(x) = (\tilde{\beta_n}(x)) + \oplus_{i=1}^{k} P$ and $g = (g_l)$ for some projection $g_l \in P$ for $l \in \mathbb{N}$. We have, then, for sufficient large l there is an order zero map $\tilde{\beta_l} \circ \phi\colon M_l \to P$ such that

(iii)
$$\begin{aligned} 1 - \tilde{\beta_l} \circ \phi(1) &= 1 - \tilde{\beta_l}(\phi(1)) \\ &= 1 - g_l + g_l - \tilde{\beta_l}(\phi(1)) \\ &= 1 - g_l + \tilde{\beta_l}(1 - \phi(1)) \ (\tilde{\beta_l}(1) = g_l) \\ &\preceq 1 - g_l + \tilde{\beta_l}(p_1) \\ &\preceq p_2 + p_1', \end{aligned}$$
where $p_1' \in \overline{g_lPg_l}$ is projection such that $\tilde{\beta_l}(p_1) \sim p_1'$. Note that since $\|\tilde{\beta_l}(p_1) - p_lg_l\|$ is very small, there are projections $p_1' \in \overline{g_lPg_l}$ and $p_1'' \in \overline{p_1Pp_1}$ such that $\tilde{\beta_l}(p_1) \sim p_1' \sim p_1''$. Since $p_2 \perp p_1''$, from [3, 1.1 Proposition] $p_2 + p_1' \preceq p_2 + p_1''$. Hence we have
$$\begin{aligned} 1 - \tilde{\beta_l} \circ \phi(1) &\preceq p_2 + p_1'' \\ &\preceq a \end{aligned}$$

(iv) For any normalized element $x \in M_n$ and $y \in F$ $\|[\tilde{\beta_n} \circ \phi(x), y]\| < 3\varepsilon$.

This implies that P is the tracially $\mathcal{Z}$-absorbing. ∎

The following is the main theorem in this note.

Theorem 5.4. Let $P \subset A$ be an inclusion of unital C*-algebra and E be a conditional expectation from A onto P with index finite type. Suppose that A is simple, separable, nuclear, $\mathcal{Z}$-absorbing and E has the tracial Rohklin property. P is $\mathcal{Z}$-absorbing.

Proof. It follows from Theorem 5.3 and Theorem 5.2. ∎

Corollary 5.5. Let A be an infinite dimensional simple separable unital C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G with the tracial Rokhlin property. Suppose that A is $\mathcal{Z}$-absorbing. Then we have

(1) ([8]) The fixed point algebra A^α and the crossed product $A \rtimes_\alpha G$ are $\mathcal{Z}$-absorbing.
(2) For any subgroup H of G the fixed point algebra A^H is $\mathcal{Z}$-absorbing.

Proof. (1): Since the canonical conditional expectation $E\colon A \to A^\alpha$ has the tracial Rokhlin property by Proposition 4.6, A^α is $\mathcal{Z}$-absorbing by Corollary 5.4.

Let $|G| = n$. Then $A \rtimes_\alpha G$ is isomorphic to $pM_n(A^\alpha)p$ for some projection $p \in M_n(A^\alpha)$. Since A^α is $\mathcal{Z}$-absorbing, $pM_n(A^\alpha)p$ is $\mathcal{Z}$-absorbing [32], hence $A \rtimes_\alpha G$ is $\mathcal{Z}$-absorbing.

(2): Since $\alpha_{|H}\colon H \to \mathrm{Aut}(A)$ has the tracial Rokhlin property by [5, Lemma 5.6], we know that A^H is $\mathcal{Z}$-absorbing by (1). ∎

6. Cuntz-equivalence for inclusions of C*-algebras

In this section we study the heredity for Cuntz equivalence for an inclusion of unital C*-algebras with the tracial Rokhlin proerty.

Let $M_\infty(A)^+$ denote the disjoint union $\cup_{n=1}^\infty M_n(A)^+$. For $a \in M_n(A)^+$ and $b \in M_m(A)^+$ set $a \oplus b = \mathrm{diag}(a, b) \in M_{n+m}(A)^+$, and write $a \preceq b$ if there is a sequence $\{x_k\}$ in $M_{m,n}(A)$ such that $x_k^* b x_k \to a$. Write $a \sim b$ if $a \preceq b$ and $b \preceq a$. Put $W(A) == M_\infty(A)^+/\sim$, and let $\langle a\rangle \in W(A)$ be the equivalence class containing a. Then $W(A)$ is a positive ordered abelian semigroup with equipped with the relations:

$$\langle a\rangle + \langle b\rangle = \langle a \oplus b\rangle, \quad \langle a\rangle \leq \langle b\rangle \Leftrightarrow a \preceq b, \quad a, b \in M_\infty(A)^+.$$

We call $W(A)$ a Cuntz semigroup.

Lemma 6.1. Let $P \subset A$ be an inclusion of unital C*-algebras with index finite type and $E\colon A \to P$ has the tracial Rokhlin property. Suppose that positive elements $a, b \in P^\infty$ satisfy $eb = be$ and $a \preceq eb$ in A^∞, where e is the Rokhlin projection for E. Then $a \preceq b$ in P^∞.

Proof. Since $a \preceq b$ in A^∞, there is a sequence $\{v_n\}_{n\in\mathbb{N}}$ in A^∞ such that $\|a - v_n^* e b v_n\| \to 0$ $(n \to \infty)$.

Let $E\colon A \to P$ be a conditional expectation of index finite type. Set $w_n = (\mathrm{Index}E)^{\frac{1}{2}}E^\infty(ev_n)$ for each $n \in \mathbb{N}$. Then, since

$$\begin{aligned} w_n^*bw_ne_P &= (\mathrm{Index}E)E^\infty(v_n^*e)bE^\infty(ev_n)e_P \\ &= (\mathrm{Index}E)E^\infty(v_n^*eb)E^\infty(ev_n)e_P \\ &= (\mathrm{Index}E)e_Pv_n^*ebe_Pev_ne_P \\ &= (\mathrm{Index}E)e_Pv_n^*bee_Pev_ne_P \\ &= e_Pv_n^*bv_ne_P \\ &= E^\infty(v_n^*bv_n)e_P, \end{aligned}$$

$w_n^*bw_n = E^\infty(v_n^*bv_n)$. Therefore,

$$\begin{aligned} \|a - w_n^*bw_n\| &= \|a - E^\infty(v_n^*ebv_n)\| \\ &= \|E^\infty(a - v_n^*ebv_n)\| \\ &\le \|a - v_n^*ebv_n\| \to 0 \ (n \to \infty). \end{aligned}$$

This implies that $a \preceq b$ in P^∞. ∎

Proposition 6.2. Let $P \subset A$ be an inclusion of unital C*-algebras with index finite type. Suppose that $E\colon A \to P$ has the tracial Rokhlin property. If two positive elements $a, b \in P$ satisfy $a \preceq b$ in A, then $a \preceq b$ in P.

Proof. Let $a, b \in P$ be positive elements such that $a \preceq b$ in A and $\varepsilon > 0$. Since for any constant $K > 0$ $a \preceq b$ is equivalent to $Ka \preceq Kb$, we may assume that a and b are contractive. When b is invertible, then $a = (a^{1/2}b^{-1/2})b(a^{1/2}b^{-1/2})^*$, and $a \preceq b$ in P. Hence we may assume that b has 0 as a spectrum.

Since $a \preceq b$ in A, there is $\delta > 0$ and $r \in A$ such that $f_\varepsilon(a) = rf_\delta(b)r^*$, where $f_\varepsilon\colon \mathbb{R}^+ \to \mathbb{R}^+$ by

$$f_\varepsilon(t) = \begin{cases} 0, & t \le \varepsilon \\ \varepsilon^{-1}(t-\varepsilon), & \varepsilon \le t \le 2\varepsilon \\ 1, & t \ge 2\varepsilon. \end{cases}$$

Set $a_0 = f_\delta(b)^{1/2}r^*rf_\delta(b)^{1/2}$. Then $f_\varepsilon(a) \sim a_0$. Set a continuos function $g_\delta(t)$ on $[0,1]$ by

$$g_\delta(t) = \begin{cases} \delta^{-1}(\delta - t) & 0 \le t \le \delta \\ 0 & \delta \le t \le 1. \end{cases}$$

Since b has 0 as a spectrum, $g_\delta(b) \neq 0$ and $g_\delta(b)f_\delta(b) = 0$. Note that $g_\delta(b)(= c)$ belongs to $\overline{bPb}$. Therefore, there are positive elements a_0 in $\overline{bAb}$ and c in $\overline{bPb}$ such that $(a-\varepsilon)_+ \preceq a_0 + c$ in A. Indeed,

$$\begin{aligned} (a-\varepsilon)_+ &\preceq f_\varepsilon(a) \\ &\sim a_0 \\ &\preceq a_0 + c. \ ([3, \text{Proposition 1.1}]) \end{aligned}$$

Take a Rokhlin projection $e \in A' \cap A^\infty$ for E. Then there is a projection $g \in P' \cap P^\infty$ such that $(1-g) \preceq ec$. Hence $(a-\varepsilon)_+(1-g) \preceq ec$ in A^∞. By Lemma 6.1 $(a-\varepsilon)_+(1-g) \preceq c$ in P^∞.

We have then in P^∞

$$\begin{aligned}(a-\varepsilon)_+ &= (a-\varepsilon)_+g + (a-\varepsilon)_+(1-g)\\ &= \beta((a-\varepsilon)_+) + (a-\varepsilon)_+(1-g)\\ &\preceq \beta(a_0) + (a-\varepsilon)_+(1-g) \ \ ([3, \text{Proposition 1.1}])\\ &\preceq \beta(a_0) + c \in \overline{bP^\infty b}, \ ([3, \text{Proposition 1.1}])\end{aligned}$$

where $\beta\colon A \to gP^\infty g$ by Lemma 4.7. Hence $(a-\varepsilon)_+ \preceq b$ in P^∞.

Since $\varepsilon > 0$, we have $a \preceq b$ in P^∞, and $a \preceq b$ in P. ∎

7. Strict comparison property

In this section we study the strictly comparison property for a Cuntz semigroup and show that for an inclusion $P \subset A$ of exact, unital C*-algebras with the tracial Rokhlin property if A has strictly comparison, then so does P. When $E\colon A \to P$ has the Rokhlin property, the statement is proved in [23].

A dimension function on a C*-algebra A is a function $d\colon M_\infty(A)^+ \to \mathbb{R}^+$ which satisfies $d(a \oplus b) = d(a) + d(b)$, and $d(a) \leq d(b)$ if $a \preceq b$ for all $a, b \in M_\infty(A)^+$. If τ is a positive trace on A, then

$$d_\tau(a) = \lim_{n\to\infty} \tau(a^{\frac{1}{n}}) = \lim_{\varepsilon\to 0+} \tau(f_\varepsilon(a)), \quad a \in M_\infty(A)^+$$

defines a dimension function on A. Every lower semicontinuous dimension function on an exact C*-algebra arises in this way ([2, Theorem II.2.2], [7], [11]). For the Cuntz semigroup $W(A)$ an additive order preserving mapping $\tilde{d}\colon W(A) \to \mathbb{R}^+$ is given by $\tilde{d}(\langle a\rangle) = d(a)$ from a dimension function d on A. We use the same symbol to denote the dimension function on A and the corresponding state on $W(A)$.

Recall that an C*-algebra A has strict comparison if whenenver $x, y \in W(A)$ are such that $d(x) < d(y)$ for all dimension function d on A, then $x \leq y$. If A is simple, exact and unital , then the strictly comparison property is equivalent to the strictly comparison proeprty by traces, that is, for all $x, y \in W(A)$ one has that $x \leq y$ if $d_\tau(x) < d_\tau(y)$ for all tracial states τ on A ([28, Corollary 4.6]).

Let $\mathrm{T}(A)$ be the set of all traces on a C*-algebra A.

Proposition 7.1. Let $E\colon A \to P$ be of index finite type and has the tracial Rokhlin property. Then the restriction map defines a bijection from the set $\mathrm{T}(A)$ to the set $\mathrm{T}(P)$.

Proof. Since $E\colon A \to P$ has the tracial Rokhlin property, for any n there are a projection $g_n \in P^\infty$ and an injective *-homomorphism β_n from A to $g_nP^\infty g_n$ such that $\beta(x) = xg_n$ for all $x \in P$ and $\tau^\infty(1-g_n) < \frac{1}{n}$ for any $\tau \in \mathrm{T(P)}$. Then we show that the map $\mathrm{T(P)} \ni \tau \mapsto \mathrm{w^*-lim}(\tau^\infty \circ \beta_n) \in \mathrm{T(A)}$ is an inverse of the restriction map, where τ^∞ be the extended tracial state of τ on P^∞ and w*− means the weak*-limit.

Let R be the restriction map of $\tau \in \mathrm{T(A)}$ to P. For $\tau \in T(P)$ $\tau^\infty \circ \beta_n$ is a tracial state on A. Then for any $a \in P$

$$\begin{aligned} R(\mathrm{w}^*-\lim(\tau^\infty \circ \beta_n))(a) &= \mathrm{w}^*-\lim(\tau^\infty(ag_n)) \\ &= \tau^\infty(a) \\ &= \tau(a). \end{aligned}$$

Hence R is surjective.

Suppose that $R(\tau_1) = R(\tau_2)$ for $\tau_1, \tau_2 \in T(A)$. Since $E\colon A \to P$ is of index finite type, there exists a quasi-basis $\{(u_i, u_i^*)\}_{i=1}^n \subset A \times A$ such that for any $x \in A$

$$x = \sum_{i=1}^n E(xu_i)u_i^* = \sum_{i=1}^n u_i E(u_i^* x).$$

Let $\tau_1 - \tau_2 = (\tau_1 - \tau_2)_+ - (\tau_1 - \tau_2)_-$ be the Jordan decomposition of $\tau_1 - \tau_2$. Since E is completely positive, $\tau \circ E$ is completely positive. We have then by the Schwarz inequality

$$\begin{aligned} |(\tau_1 - \tau_2)_+(E(xu_i)u_i^*)\tau(E(xu_i)u_i^*)^*| &\le |(\tau_1 - \tau_2)_+(E(xu_i)u_i^*(E(xu_i)^* u_i^*)^*)| \\ &= |(\tau_1 - \tau_2)_+(E(xu_i)u_i^* u_i E(xu_i)^*)| \\ &\le \|u_i\|^2 |(\tau_1 - \tau_2)_+(E(xu_i)E(xu_i)^*)| = 0. \end{aligned}$$

Hence $(\tau_1 - \tau_2)_+(E(xu_i)u_i^*) = 0$. This means that

$$(\tau_1 - \tau_2)_+(x) = \sum_{i=1}^n (\tau_1 - \tau_2)_+(E(xu_i)u_i^*) = 0.$$

Hence $(\tau_1 - \tau_2)_+ = 0$. Similarly, $(\tau_1 - \tau_2)_- = 0$. Hence, $\tau_1 - \tau_2 = 0$, and R is injective. ∎

Theorem 7.2. Let $P \subset A$ be an inclusion of unital C*-algebras with index finite type. Suppose that A is simple and exact and A has strict comparison, and $E\colon A \to P$ has the tracial Rokhlin property. Then P has strict comparison.

Proof. Since $E\colon A \to P$ is of index finite type and A is simple, P is exact and simple. Note that the strict comparison property is equivalent to the the strict comparison property given by traces, i.e., for all $x, y \in \mathcal{C}(A)$ one has that $x \le y$ if $d_\tau(x) < d_\tau(y)$ for all tracial states τ on A (see [29, Remark 6.2] [28]).

Since $E \otimes id\colon A \otimes M_n \to P \otimes M_n$ is of index finite type and has the tracial Rokhlin property, it suffices to verify the condition that whenever $a, b \in P$ are positive elements such that $d_\tau(a) < d_\tau(b)$ for all $\tau \in \mathrm{T(P)}$, then $a \preceq b$.

Let $a, b \in P$ be positive elements such that $d_\tau(a) < d_\tau(b)$ for all $\tau \in \mathrm{T(P)}$. Then by Proposition 7.1 we have $d_\tau(a) < d_\tau(b)$ for all tracial states $\tau \in \mathrm{T}(A)$. Since A has strict comparison, $x \preceq y$ in A. Therefore By Proposition 6.2 $a \preceq b$ in P, and P has strict comparison. ∎

Corollary 7.3. Let $P \subset A$ be an inclusion of unital C*-algebras with index finite type. Suppose that A is simple and the order on projections on A is determined by traces, and $E\colon A \to P$ has the tracial Rokhlin property. Then the order on projections on P is determined by traces.

Proof. Since $E \otimes id\colon A \otimes M_n \to P \otimes M_n$ is of index finite type and has the tracial Rokhlin property, it suffices to verify the condition that whenever $p, q \in P$ are projections such that $\tau(p) < \tau(q)$ for all $\tau \in \mathrm{T(P)}$, then p is the Murray-von Neumann equivalent to a subprojection of q in P.

Let $p, q \in P$ be projections such that $\tau(p) < \tau(q)$ for all $\tau \in \mathrm{T(P)}$. Then by Proposition 7.1 we have $\tau(p) < \tau(q)$ for all tracial states $\tau \in \mathrm{T}(A)$. Since the order on projections on A, p is the Murray-von Neumann subequivalnt to q in A, and $p \preceq q$ in A by [26, Proposition 2.1]. Therefore By Proposition 6.2 $p \preceq q$ in P, and p is the Murray-von Neumann equivalent to a subprojection of q in P. Hence the order on projections on P is determined by traces. ∎

The following is well-known.

Lemma 7.4. let A be an exact C*-algebra and p be a projection of A. Suppose that A has strictly comparison. Then so does pAp.

Proof. Since for each $n \in \mathbb{N}$ $M_n(A)$ is exact and has strictly comparison, we have only to show that whenever $a, b \in pAp$ are positive elements such that $d_\tau(a) < d_\tau(b)$ for all tracial states $\tau \in \mathrm{T}(pAp)$, then $a \preceq b$.

Let $a, b \in pAp$ be positive elements such that $d_\tau(a) < d_\tau(b)$ for all tracial states $\tau \in \mathrm{T}(pAp)$. Then for any tracial $\rho \in \mathrm{T}(A)$ the restriction $\rho_{|pAp}$ belongs to $\mathrm{T}(pAp)$. Since $d_\rho(a) = d_{\rho_{|pAp}}(a) < d_{\rho_{|pAp}}(b) = d_\rho(b)$, we have $a \preceq b$ in A by the assumption. Hence there is a sequence $\{x_n\} \subset A$ such that $\|x_n^* b x_n - a\| \to 0$. Set $y_n = p x_n p \in pAp$ for each $n \in \mathbb{N}$, then

$$\begin{aligned} \|y_n^* b y_n - a\| &= \|p x_n^* p b p x_n p - pap\| \\ &= \|p x_n^* b x_n p - pap\| \\ &\leq \|x_n^* b x_n - a\| \to 0, \end{aligned}$$

and $a \preceq b$ in pAp. Therefore, pAp has strict comparison. ∎

Corollary 7.5. Let A be an infinite dimensional simple separable unital C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G with the tracial Rokhlin property. Suppose that A is exact and has strict comparsion. Then we have

(1) [8] The fixed point algebra A^α and the crossed product $A \rtimes_\alpha G$ have strict comparioson.
(2) For any subgroup H of G the fixed point algebra A^H has strict comparison.

Proof. (1) Since the canonical conditional expectation $E\colon A \to A^\alpha$ has the tracial Rokhlin property by Proposition 4.6, A^α has strict comparison by Theorem 7.3.

Let $|G| = n$. Then $A \rtimes_\alpha G$ is isomorphic to $pM_n(A^\alpha)p$ for some projection $p \in M_n(A^\alpha)$. Since A^α has strict comparison, $pM_n(A^\alpha)p$ has strict comparison by Lemma 7.4, hence $A \rtimes_\alpha G$ has strict comparison.

(2): Since $\alpha_{|H}\colon H \to \mathrm{Aut}(A)$ has the tracial Rokhlin property by [5, Lemma 5.6], we know that A^H has strict comparison by (1). ∎

Similarly we have the following.

Corollary 7.6. Let A be an infinite dimensional simple separable unital C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G with the tracial Rokhlin

property. Suppose that the order on projections on A is determined by traces. Then we have

(1) [8] The order on projections on the fixed point algebra A^α and the crossed product $A \rtimes_\alpha G$ is determined by traces.
(2) For any subgroup H of G the order on projections on the fixed point algebra A^H is determined by traces.

Acknowledgement The authors thank Professor Yasuhiko Sato for fruitful discussion, in particlular, about Lemma 4.10.

References

[1] D. Archey, *Crossed product C*-algebras by finite group actions with the tracial Rokhlin property*, arXiv:0902.2865.
[2] B. Blackadar and D. Handelman, *Dimension functions and traces on C*-algebras*, J. Funct. Anal. **45**(1982), 297 - 340.
[3] J. Cuntz, *Dimension functions on simple C*-algebras*, Math. Ann. 233(1978), 145 - 153.
[4] G. A. Elliott and Z. Niu, *On tracial approximation*, Journal of Functional Analysis **254**(2008), 396 - 440.
[5] S. Echterhoff, W. Luck, N. C. Phillips, S. Walters, *The structure of crossed products of irrational rotation algebras by finite subgroups of $SL_2(\mathbb{Z})$* J. Reine Angew. Math. **639**(2010), 173–221.
[6] Q. Fan and X. Fang, *Stable rank one and real rank zero for crossed products by finite group actions with the tracial Rokhlin property*, Chinese Anal. Math. **30**(2009), 179 - 186.
[7] U. Haagerup, *Every quasi-trace on an exact C*-algebra is a trace*, preprint (1991).
[8] I. Hirshberg and J. Orovitz, *Tracially $\mathcal{Z}$-absorbing C*-algebras*, J. Funct. Anal. **265**(2013), 765-785.
[9] M. Izumi, *Finite group actions on C*-algebras with the Rohlin property–I*, Duke Math. J. **122**(2004), p. 233–280.
[10] X. Jiang and H. Sue, *On a simple unital projectionless C*-algebras* Amer. J. Math **121**(1999), p. 359–413.
[11] E. Kirchberg, *On the existence of traces on exact stably projectionless simple C*-algebras*, in Operator Algebras and their Applications, eds. P. A. Fillmore and J. A. Mingo, Fields Institute Communications, Vol. 13 (American Mathematical Society, 1995), pp. 171–172.
[12] A. Kishimoto, *Outer automorphisms and reduced crossed products of simple C*-algebras*, Commun. Math. Phys. **81**(1981), 429 - 435..
[13] K. Kodaka, H. Osaka, and T. Teruya, *The Rohlin property for inclusions of C*-algebras with a finite Watatani Index*, Operator structures and Dynamical Systems, Contemporary Mathematics 503(2009), 179 - 195.
[14] H. Lin, *Tracial AF C*-algebras*, Trans. Amer. Math. Soc. 353(2001), 693–722.
[15] H. Lin, *An Introduction to the Classification of Amenable C*-algebras*, World Scientific, River Edge NJ, 2001.
[16] H. Lin, *Simple nuclear C*-algebras of tracial topological rank one*, J. Funct. Anal. 251(2007), 601–679, arXiv:math.OA/0401240, 2004.
[17] T. A. Loring, *Lifting Solutions to Perturbing Problems in C*-algebras*, Fields Institute Monographs no. 8, American Mathematical Society, Providence RI, 1997.
[18] H. Matui and Y. Sato, *Strict comparison and Z -absorption of nuclear C*-algebras*, Acta Math. **209**(2012), no. 1, 179–196.
[19] H. Osaka, *SP-Property for a pair of C*-algebras*, J. Operator Theory **46**(2001), 159 - 171.
[20] H. Osaka and N. C. Phillips, *Crossed products by finite group actions with the Rokhlin property*, Math. Z. 270(2012), 19–42, arXiv:math.OA/0704.3651.
[21] H. Osaka and N. C. Phillips, in preparation.
[22] H. Osaka and T. Teruya, *Strongly self-absorbing property for inclusions of C*-algebras with a finite Watatani index*, Trans. Amer. Math. Soc. **366**(2014) no. 3 1685–1702.
[23] H. Osaka and T. Teruya, *Nuclear dimension and pureness for an inclusion of unital C*-algebras*, preprint, arXiv:1111.1808.

[24] N. C. Phillips, *The tracial Rokhlin property for actions of finite groups on C*-algebras* Amer. J. Math. 133(2011), no. 3, 581–636, arXiv:math.OA/0609782.
[25] M. Rordam, *On the structure of simple C*-Algebras tensored with a UHF-algebra*, J. Funct. Anal. **100**(1991), 1 - 17.
[26] M. Rordam, *On the structure of simple C*-Algebras tensored with a UHF-algebra II*, J. Funct. Anal. **107**(1992), 255 - 269.
[27] M. Rørdam, *Classification of nuclear, simple C*-algebras*, Encyclopaedia Math. Sci. , 126, Springer, Berlin, 2002.
[28] M. Rordam, *The stable and the real rank of $\mathcal{Z}$-absorbing C*-algebras*, Inter. J. Math. **10**(2004), 1065–1084.
[29] M. Rørdam and W. Winter, *The Jiang-Su algebra revisited*, J. reine angew. Math. **642**(2010), 129 - 155.
[30] Y. Sato, private communication.
[31] A. S. Toms, *Characterizing classifiable AH algebras*, C. R. Math. Acad. Sci. Soc. R. Can. **33**(2011), no. 4, 123–126.
[32] A. S. Toms and W. Winter, *Strongly self-absorbing C*-algebras*, Trans. Amer. Math. Soc. **359**(2007), 3999 - 4029.
[33] Y. Watatani, *Index for C^*-subalgebras*, Mem. Amer. Math. Soc. **424**, Amer. Math. Soc., Providence, R. I., (1990).
[34] W. Winter, *Covering dimension for nuclear C*-algebras I*, J. Funct. Anal. 199(2003), 535-556.
[35] W. Winter and J. Zacharias, *The nuclear dimension of C*-algebras*, Adv. Math. **224**(2010), 461–498.
[36] X. Yang and X. Fang, *The tracial class property for crossed products by finite group actions*, Abstract and Applied Analysis 2012, Article ID 745369.

Department of Mathematical Sciences, Ritsumeikan University, Kusatsu, Shiga, 525-8577 Japan
E-mail address: osaka@se.ritsumei.ac.jp

Faculty of Education, Gunma University, 4-2 Aramaki-machi, Maebashi City, Gunma, 371-8510, Japan
E-mail address: teruya@gunma-u.ac.jp